

Subject: Electric Tumor Treatment Field (TTF)
Guideline #: CG-DME-44
Status: Revised

Publish Date: 11/17/2022
Last Review Date: 11/10/2022

Description

This document addresses electrical fields known as “tumor treatment fields (TTF)” that are created by low-intensity, intermediate frequency (100–200 kilohertz [kHz]) electric currents delivered to the malignant tumor site by insulated electrodes placed on the skin surface. TTF are felt to cause tumor cell death (apoptosis) by disrupting the assembly of microtubules during later stages of cell division.

Clinical Indications

Medically Necessary:

The use of FDA approved devices to generate electric tumor treatment fields (TTF) to treat histologically-confirmed supratentorial glioblastoma (known also as glioblastoma multiforme [GBM] or World Health Organization [WHO] grade IV astrocytoma) is considered **medically necessary** when criteria A or B are met:

- A. As an adjunctive treatment in newly diagnosed GBM when **all** the following criteria are met:
1. Initial treatment with debulking surgery or biopsy followed by chemoradiation with concomitant temozolomide and radiotherapy has been completed with no documented tumor progression*; **and**
 2. TTF is used in combination with temozolomide; **and**
 3. TTF is initiated within 7 weeks from final dose of temozolomide and radiotherapy; **and**
 4. Individual has Karnofsky Performance Status (KPS) score of 70 or higher **or** Eastern Cooperative Oncology Group (ECOG) performance status 0-1; **and**
 5. Individual or caregiver has been trained and is willing and able to apply and use the device at least 18 hours, per day on average.
- B. In recurrent GBM when **all** the following criteria are met:
1. TTF is used as a monotherapy; **and**
 2. Individual has KPS score of 70 or higher or ECOG performance status 0-1; **and**
 3. Individual or caregiver has been trained and is willing and able to apply and use the device at least 18 hours, per day, on average.

Continuation of TTF therapy beyond the initial 90 days is considered **medically necessary** when *both* of the following criteria are met:

- Documentation of compliant use must be reported every 90 days as evidenced by the device monitor report showing the individual is using the device at least 18 hours per day, on average; **and**
- There is no documented tumor progression*.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

*See [discussion section](#) regarding *tumor progression* criteria

Not Medically Necessary:

The use of devices to generate electric tumor treatment fields (TTF) is considered **not medically necessary** when the criteria above are not met and for all other malignant tumors.

The use of enhanced computer treatment planning software (such as NovoTal) is considered **not medically necessary** in all cases.

Coding

The following codes for treatments and procedures applicable to this guideline are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

When services may be Medically Necessary when criteria are met:

CPT

77299 Unlisted procedure, therapeutic radiology clinical treatment planning [when specified as plan for using a medically necessary electrical stimulation device for TTF for GBM]

HCPCS

A4555 Electrode/transducer for use with electrical stimulation device used for cancer treatment, replacement only
 E0766 Electrical stimulation device used for cancer treatment, includes all accessories, any type

ICD-10 Diagnosis

C71.0-C71.9 Malignant neoplasm of brain
 Z85.841 Personal history of malignant neoplasm of brain

When services are Not Medically Necessary:

For the procedure and diagnosis codes listed above when criteria are not met, and for all other diagnoses not listed.

When services are also Not Medically Necessary:

CPT

77299 Unlisted procedure, therapeutic radiology clinical treatment planning [when specified as treatment planning for use of an electrical stimulation device for TTF using enhanced computer software (e.g., NovoTal)]

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

ICD-10 Diagnosis

All diagnoses

Discussion/General Information***Glioblastoma Multiforme (GBM)***

Glioblastoma (WHO grade IV astrocytoma), also known as GBM (National Cancer Institute [NCI], 2022), has a peak incidence between the ages of 45 and 70 years. GBM is the most frequently occurring brain tumor accounting for approximately 12% to 15% of all brain tumors and 50% to 60% of all astrocytic tumors. Giant cell glioblastoma and gliosarcoma are two histologic variants of GBM. The 5-year survival rate for GBM is between 1% and 19%, depending upon age. Treatment of GBM is challenging due to the inability of most systemic therapy agents to cross the blood-brain barrier and the propensity of high-grade gliomas to recur (National Comprehensive Cancer Network® [NCCN], V2.2022). The delivery of low-intensity, intermediate-frequency alternating electric fields has been demonstrated to have an inhibitory effect on proliferating cells. Exposure to the alternating fields lead to mitotic disruption and ultimately mitotic cell death. Nonproliferating cells remain unaffected by these fields (Fabian, 2019).

NovoTTF™-100A System (NovoCure™ Ltd., Portsmouth, NH; Haifa, Israel) received U.S. Food and Drug Administration (FDA) premarket approval (PMA) in 2011. The device is now marketed as Optune® (NovoCure Ltd., Portsmouth, NH, Haifa, Israel). Optune, a portable, non-invasive device designed for the delivery of TTF to the head, should be worn at least 18 hours a day in order to obtain the best treatment response. The device was originally approved as a novel device to treat adults aged 22 years or older with GBM that recurs or progresses after receiving chemotherapy and radiation therapy. On October 5, 2015, the FDA approved the use of Optune in combination with temozolomide for the treatment of adults with newly diagnosed, supratentorial GBM following maximal debulking surgery and radiation therapy.

The first generation Optune system weighed approximately 6 pounds, the weight of the second generation Optune system has been reduced to 2.7 pounds. The second-generation device has also been redesigned to generate significantly less noise. These improvements may result in increased compliance. In a study by Kinzel and associates (2019), 10 individuals with GBM who were currently using the first generation TTF device, were transitioned to the second-generation device. Compliance improved in 4 users, was maintained in 5 users and decreased in 1 user. A total of 3/10 users had not reached 75% compliance level with the first-generation device but did achieve ≥75% compliance following transition to the second-generation device.

Newly diagnosed GBM

Current standard treatment for newly diagnosed GBM consists of tumor resection followed by daily low dose temozolomide administered concurrently with external beam radiotherapy followed by adjuvant temozolomide with alternating electric field therapy (NCCN, 2022). Radiochemotherapy is followed by adjuvant temozolomide given for 6 to 12 months. The prognosis for individuals with GBM is poor, with a 1-year survival rate of less than 40%.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Stupp and colleagues (2015) evaluated the safety and efficacy of TTF in individuals with newly diagnosed GBM following chemoradiation therapy. In a multi-center clinical trial, 695 individuals were randomized (2:1) to either TTF therapy (worn at least 18 hours/day) with temozolomide or temozolomide alone. The primary endpoint was identified as progression-free survival (PFS) time in the intent-to-treat (ITT) population (significant threshold, $p \leq 0.01$).

An interim analysis conducted on the first 315 participants who had completed at least 18 months of follow-up revealed median PFS in the TTF plus temozolomide group of 7.1 months (95% confidence interval [CI], 5.9-8.2 months) compared to 4.0 months (95% CI, 3.3-5.2 months) in the temozolomide group (Hazard Ratio [HR] 0.62; 98.7% CI, 0.43-0.89; stratified log-rank, $p=0.001$). Median overall survival (OS, secondary endpoint) in the per-protocol population was 20.5 months (95% CI, 16.7-25.0 months) in the TTF plus temozolomide group versus 15.6 months (95% CI, 13.3-19.1 months) (HR, 0.64; 99.4% CI, 0.42-0.98; $p=0.004$) in the temozolomide alone group. Based on the interim analysis results, the study was terminated and individuals in the control group were offered TTF therapy in addition to temozolomide. A total of 11 individuals crossed over and began using TTF therapy. With the exception of a higher incidence of localized skin reactions in the TTF plus temozolomide group, the incidence, distribution, and severity of adverse events (AEs) were similar across both treatment groups. The final analysis of the data was consistent with the interim analysis results (Stupp, 2017).

Results from an industry-sponsored pilot study of TTF therapy alone and TTF therapy in combination with chemotherapy for individuals with diagnosed GBM were reported by Kirson and associates (2009b). In this single arm study, the first group included 10 individuals with recurrent GBM after failure of maintenance temozolomide, and 10 individuals with newly diagnosed GBM treated with TTF therapy combined with temozolomide were in the second cohort. All 20 individuals were treated for an average of 1 year (range 2.5-24 months) continuously. The first group was compared to a matched group of 18 concurrent controls who received salvage chemotherapy for relapsed/recurrent GBM. The TTF-chemotherapy group was compared to a matched group of 32 concurrent controls who received temozolomide alone. In addition, OS for both cohorts was compared to matched historical control data. Data for the first group were reported by Kirson and colleagues in 2007. For the group of 10 individuals with newly diagnosed GBM, PFS was significantly different (HR 3.32; 95% CI, 1.9-5.9; $p=0.0002$) between the TTF-chemotherapy group compared to the matched concurrent and historical controls. The difference in OS was also significant ($p=0.0018$). The authors concluded TTF therapy may also be an effective sensitizer when used concurrently with chemotherapeutic agents.

The use of adjuvant alternating electric field therapy when used as an initial therapy along with temozolomide for individuals with anaplastic gliomas/ glioblastoma with good performance status following standard radiotherapy and concurrent temozolomide is a NCCN category 1 recommendation.

Recurrent or Progressive GBM

Stupp and associates (2012) conducted a phase III, pivotal, multinational, randomized controlled trial (RCT) upon which the initial PMA was based. Between September 2006 and May 2009, 28 clinical centers enrolled 237 adult participants with relapsed or progressive GBM despite conventional therapy (e.g., surgery and chemo-radiotherapy followed by chemotherapy). A total of 120 participants were randomized in a 1:1 ratio to receive monotherapy with

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

TTF therapy and 117 participants were randomized to the group treated with available best standard care (BSC) chemotherapies as practiced at each of the participating clinical centers. Chemotherapy agents considered as BSC during the trial included platinum-based chemotherapy (i.e., carboplatin); nitrosureas (BCNU); procarbazine; combination of procarbazine, lomustine and vincristine (PCV); temozolomide; and bevacizumab. A period of 28 days of treatment with the TTF device was considered one full treatment course. Participants treated with the TTF device were allowed to take breaks from treatment up to an hour, twice per day for personal needs such as showers. The primary endpoint of the study was OS. Secondary endpoints included PFS at 6 months (PFS6), time-to-progression (TTP), 1-year survival rate, quality of life (QOL), and radiological response. Participants were seen in clinic monthly, and magnetic resonance imaging (MRI) was performed after 2, 4, and 6 months from initiation of treatment and subsequent MRIs were done according to local practice until disease progression. Medical follow-up continued for 2 months after disease progression. Monthly telephone interviews with the participants' caregivers were used to assess participant mortality rates.

A total of 97% (116) of 120 enrollees in the TTF group started treatment and 93 participants (78%) completed 1 cycle (4 weeks) of therapy. Discontinuation of TTF therapy occurred in 27 participants due to noncompliance or the inability to handle the device. For each TTF treatment month, the median compliance was 86% (range 41-98%), which equaled a mean use of 20.6 hours per day. In the BSC (active control) group, 113 (97%) of the 117 assigned participants received chemotherapy, 112/113 participants completed one full treatment course. In the BSC cohort, 21 participants did not return to the site and details on disease progression and toxicity were not available. The median survival of 6.6 months in the TTF therapy group was marginally higher than 6 months in the BSC group (HR 0.86; 95% CI, 0.66-1.12; p=0.27). For both groups, 1-year survival was 20%. The survival rates for 2 and 3 years were 8% (95% CI: 4, 13) and 4% (95% CI: 1, 8) versus 5% (95% CI: 3, 10) and 1% (95% CI: 0, 3) for the TTF cohort compared to the BSC cohort, respectively. With a median follow-up of 39 months, 93% (220 participants) had died. Objective radiological responses (partial response [PR] and complete response [CR]) were noted in 14 participants in the TTF group and 7 in the BSC group, with a calculated response rate of 14.0% (95% CI, 7.9-22.4%) compared to 9.6% (95% CI, 3.9-18.8%), respectively. Individuals in the TTF therapy group reported localized reactions which resolved with topical steroids. BSC participants experienced grade 2-4 events by organ system related to the pharmacologic activity of chemotherapy agents utilized. This trial represents the first phase III clinical trial of the TTF device using the first-generation device. The primary endpoint of the trial, improved OS compared to chemotherapy, was not reached. The TTF group did demonstrate similar efficacy to chemotherapy regimens with a lower toxicity profile and better quality of life (Fabian, 2019).

In 2014, Mrugala and associates analyzed the registry data of all individuals with recurrent GBM who had undergone therapy with the TTF device in 91 cancer centers (n=457). The median survival rate was 9.6 months. This was significantly longer than the 6.6 months reported in the Stupp (2012) trial. The individuals included in the registry data analysis also reported more than double the overall 1- and 2-year survival rates than reported in the Stupp 2012 trial. This may have been associated with the number of recurrences experienced prior to the application of the TTF device. Individuals in the data registry cohort were more likely to experiencing their first recurrence (33%) compared to individuals in the Stupp (2012) trial (9%). Compliance is also a prognostic factor of OS. Individuals with 75% or greater compliance, defined as daily compliance $\geq 75\%$ or ≥ 18 hours daily, had a median OS of 13.5 months compared to 4.0 months in those with suboptimal compliance.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Electric Tumor Treatment Field (TTF)

Vymazal and colleagues (2015) analyzed the response patterns in individuals with recurrent GBM who exhibited an objective response in two previous studies in order to evaluate the baseline characteristics of those individuals who responded and to evaluate the relationship between compliance with use and efficacy outcomes. The analysis was completed on one pilot study (n=10) and a phase III trial (n=237) in which TTF therapy was compared to standard chemotherapy. Between both studies, TTF was administered as monotherapy in 130 individuals. Across both trials, there was a 15% response rate (16/110 with a 4% CR rate). There were no significant differences in baseline characteristics between the responder and non-responder groups. In those in which a response was noted, there was frequently a delayed response; the tumor would initially continue to grow before responding to treatment. Analysis supported that an increase in compliance was associated with better treatment response and longer OS. The extent of treatment response in those who exhibited a response was significantly dependent on compliance.

Treatment recommendations for brain tumors include surgical resection, radiation therapy and/or chemotherapy as treatment options. There is no established second-line therapy for recurrent gliomas (NCI, 2022; NCCN, V2.2022). The NCCN panel designates a 2B recommendation for alternating electric field therapy in the treatment of recurrent GBM. The evidence supports that the use of TTF in recurrent GBM is associated with improved OS when used consistently with a trend towards higher levels of survival associated with increasing compliance (Toms, 2019).

Tumor progression ([Return to Clinical Indications](#))

Tumor progression, based on Macdonald criteria was defined as at least one of the following:

- Tumor growth of greater than 25% of the product of 2 perpendicular diameters compared to the smallest tumor area measured
- Appearance of one or more new tumors in the brain (diagnosed radiologically as GBM).

Compliance with therapy

The Optune device is designed to be used at least 18 hours every day in order to maximize treatment benefit. Continuous application is needed to reach and maintain therapeutic efficacy (Regev, 2021). The compliance rate, the number of days in which the minimum 18 hours daily use is reached, is an independent predictor of overall survival (Toms, 2019). In a systematic review and meta-analysis (Regev, 2021) reported on the pooled effect of compliance on OS rates of those with a TTF use daily compliance of 75% or more (10.3 months) compared to those with a TTF daily compliance of less than 75% (5.7 months). The 6-, 9-, and 12-month OS rates of those who used the TTF device 75% or more improved compared to those who used the TTF device less than 75% (73.5% [95% CI = 67.5-80], 54.3% [95% CI = 47.7-61.9], and 44.4% [95% CI = 37.9-51.9] compared to 48% [95% CI = 41-56.2], 40.6% [95% CI = 33.7-49], and 32.5% [95% CI = 25.8-40.9], respectively). An internal log file monitors compliance with therapy.

Malignant Pleural Mesothelioma

Pleural mesothelioma cancer develops in the mesothelial surface of the lungs and is primarily associated with asbestos exposure. Malignant mesothelioma is rare in the United States, with approximately 2500 new cases

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

diagnosed each year. Pleural mesothelioma accounts for more than 75% of mesothelioma cases. Most individuals present with advanced disease limiting treatment options available. Median OS is 1 year with a 5-year OS of about 10% (NCCN, 2022).

In May 2019, the FDA approved the use of a modified version of the TTF device (Optune Lua™ previously known as NovoTTF™-100L) for the first-line treatment of adults with unresectable, locally advanced or metastatic malignant pleural mesothelioma to be used concurrently with pemetrexed and platinum-based chemotherapy. The device was approved under the Humanitarian Device Exemption (HDE) and identifies the specific population that the modified device is intended to treat. The HDE was not reviewed by the FDA advisory panel, the approval is based upon the previous review and approval of the similar GBM device. The Optune Lua operates on the same principle as the device used to treat GBM but includes different technological characteristics and area of application.

Ceresoli and colleagues (2019) evaluated the activity of TTF therapy used in combination with systemic chemotherapy in treating Stage IV unresectable malignant pleural mesothelioma (STELLAR study). Participants in the Phase II, prospective, single arm study (n=80) were treated with pemetrexed and cisplatin or carboplatin in combination with TTF therapy to the thorax until radiological disease progression or unacceptable toxicity was seen. The primary endpoint was OS time from diagnosis until the date of death. The median OS was 18.2 months (95% CI, 12.1-25.8). The 1-year survival rate was 62% and the 2-year survival rate was 42%. This OS was compared to the OS reported in two recent randomized trials involving malignant pleural mesothelioma chemotherapy regimens. These studies evaluated the addition of bevacizumab or nintedanib to a standard cisplatin and pemetrexed regimen. Both studies reported increased OS in the study groups compared to control groups, 18.8 months (95% CI, 15.9-2.6) and 18.3 months (15.2-28.8) compared to 16.1 months (14.0-17.9) and 14.2 months (95% CI 12.3-20.9) respectively. A total of 32 individuals (40%) reported severe AEs during the study, with anemia and neutropenia being unrelated to device use. Device-related AEs primarily consisted of skin reactions and were reported in 66% of the individuals, with 5% of these reactions being severe enough to result in treatment interruption. Randomized trials which compare standard chemotherapy with and without concurrent TTF therapy are needed to further evaluate any potential incremental benefit of this therapy over the current standard of care.

In summary, the available evidence regarding the use of TTF in treating stage IV non-curative mesothelioma in combination with standard chemotherapy does not demonstrate that use provides cost effective, therapeutically equivalent outcomes over the use of standard chemotherapy therapy alone.

Other Solid Tumors

In addition to TTF treatment for GBM, phase III trials are underway in other types of malignancies, including brain metastasis, non-small cell lung cancer (NSCLC), ovarian cancer and pancreatic carcinoma. The FDA has granted breakthrough device designation to the NovoTTF-200T for liver cancer. This designation was granted based upon the unpublished results of a phase 2 study evaluating the safety and efficacy of the TTF device when used with sorafenib to treat advanced liver cancer. At this time, there are no studies which support the use of tumor treating fields for conditions other than GBM. The NCCN clinical practice guidelines do not include any recommendations regarding the use of electric TTF treatment for any condition other than GBM.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Treatment Planning Software

In 2013, the FDA approved NovoTal through a PreMarket Approval (PMA) supplement. NovoTal is an algorithmic software package which allows treating physicians, who have completed a certification program, to create individualized treatment maps. The standard treatment plan developed by the manufacturer uses post-contrast MRI sequences to develop a treatment plan. Treating physicians using NovoTal are able to incorporate additional imaging data and other clinical considerations into TTF treatment planning (Connelly, 2016). There is a paucity of literature reporting on planning approaches in TTF treatment and their effect on clinical outcomes. Connelly and colleagues (2016) reported on the use of NovoTal in a case series of 8 individuals with grades 2-4 GBMs. In addition to contrast enhancing MRI imaging, other clinical considerations, such as the heterogeneity in contrast enhancement in tumors, were taken into account during the planning process. The authors discuss the use of alternative MRI sequences during the planning stage of treatment, but do not report on clinical outcomes, noting:

this case series demonstrates that treatment planning beyond the extent of contrast enhancement is clinically feasible and should be prospectively compared to standard treatment planning in a clinical trial setting, in order to determine the impact on patient outcomes.

Chaudhry and associates (2015) compared physician performance using the NovoTal system to conduct transducer array layout mapping to the mapping laid out by the Novocure in-house clinical team. Neuro-oncologists, medical oncologists and neurosurgeons (n=14) evaluated 5 cases of recurrent GBM and developed treatment plans. While the study demonstrated a high level of concordance in transducer array layout planning between NovoTal certified physicians and the Novocure in-house clinical team, the study did not address whether clinical outcomes were affected. The evidence does not support that the use of enhanced treatment planning software is considered effective in the use of TTF treatment.

Definitions

Cytokinesis: The cytoplasmic changes accompanying mitosis. The cleavage of the cytoplasm into daughter cells following nuclear division.

Eastern Cooperative Oncology Group (ECOG) Performance Status: A scale used to determine the individual's level of functioning. This scale may also be referred to as the WHO or Zubrod score which is based on the following scale:

- 0 Fully active, able to carry on all pre-disease performance without restriction
- 1 Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g., light house work, office work
- 2 Ambulatory and capable of all self-care but unable to carry out any work activities. Up and about more than 50% of waking hours
- 3 Capable of only limited self-care, confined to bed or chair more than 50% of waking hours
- 4 Completely disabled. Cannot carry on any self-care. Totally confined to bed or chair
- 5 Dead

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Glioblastoma multiforme: Stage IV glioblastoma, which includes WHO recognized variants, giant cell glioblastoma and gliosarcoma.

Karnofsky Performance Status Score: A 10-point scale used by healthcare providers to quickly evaluate how an individual is feeling on any given day.

100	Able to work. Normal; No complaints; No evidence of disease.
90	Able to work. Able to carry on normal activity; Minor symptoms.
80	Able to work. Normal activity with effort; Some symptoms.
70	Independent; not able to work. Cares for self; Unable to carry on normal activity.
60	Disabled; dependent. Requires occasional assistance; cares for most needs.
50	Moderately disabled; dependent. Requires considerable assistance and frequent care.
40	Severely disabled; dependent. Requires special care and assistance.
30	Severely disabled. Hospitalized, death not imminent.
20	Very sick. Active supportive treatment needed.
10	Moribund. Fatal processes are rapidly progressing

Macdonald criteria for disease progression is defined as *at least one* of the following:

- Tumor growth of greater than 25% of the product of 2 perpendicular diameters compared to the smallest tumor area measured
- Appearance of one or more new tumors in the brain (diagnosed radiologically as GBM).

Mitosis: The process by which a single parent cell divides to make two new daughter cells. Each daughter cell receives a complete set of chromosomes from the parent cell, allowing the body to grow and replace cells.

Progressive disease: Disease that is growing, spreading or getting worse.

Recurrent disease or recurrence: Disease that has recurred (come back), usually after a period of time during which the disease could not be detected. In the case of cancer, the disease may come back to the same place as the original (primary) tumor or to another place in the body (metastatic).

References

Peer Reviewed Publications:

1. Burri SH, Gondi V, Brown PD, Mehta MP. The evolving role of tumor treating fields in managing glioblastoma: guide for oncologists. *Am J Clin Oncol*. 2018; 41(2):191-196.
2. Ceresoli GL, Aerts JG, Dziadziuszko R, et al. Tumour Treating Fields in combination with pemetrexed and cisplatin or carboplatin as first-line treatment for unresectable malignant pleural mesothelioma (STELLAR): a multicentre, single-arm phase 2 trial. *Lancet Oncol*. 2019; 20(12):1702-1709.
3. Chaudhry A, Benson L, Varshaver M, et al. NovoTTF™ - 100A System (Tumor Treating Fields) transducer array layout planning for glioblastoma: a NovoTAL™ System user study. *World J Surg Oncol*. 2015; 13:316.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Electric Tumor Treatment Field (TTF)

4. Connelly J, Hormigo A, Mohilie N, et al. Planning TTFIELDS treatment using the NovoTAL system-clinical case series beyond the use of MRI contrast enhancement. *BMC Cancer*. 2016; 16(1):842.
5. Domingo-Musibay E, Galanis E. What next for newly diagnosed glioblastoma? *Future Oncol*. 2015; 11(24):3273-3283.
6. Fabian D, Guillermo Prieto Eibl MDP, Alnahhas I, et al. Treatment of glioblastoma (GBM) with the addition of tumor-treating fields (TTF): A review. *Cancers (Basel)*. 2019; 11(2):174.
7. Kanner AA, Wong ET, Villano JL, Ram Z.; EF-11 Investigators. Post hoc analysis of intention-to-treat population in phase 3 comparison of NovoTTF-100A™ System versus best physician's choice chemotherapy. *Semin Oncol*. 2014; 41(Suppl 6):S25-S34.
8. Kinzel A, Ambrogi M, Varshaver M, Kirson ED. Tumor treating fields for glioblastoma treatment: Patient satisfaction and compliance with the second-generation Optune® System. *Clin Med Insights Oncol*. 2019; 13:1179554918825449.
9. Kirson ED, Dbaly V, Tovarys F, et al. Alternating electric fields arrest cell proliferation in animal tumor models and human brain tumors. *Proc Natl Acad Sci USA*. 2007; 104(24):10152-10157.
10. Kirson ED, Giladi M, Gurvich Z, et al. Alternating electric fields (TTFIELDS) inhibit metastatic spread of solid tumors to the lungs. *Clin Exp Metastasis*. 2009a; 26(7):633-640.
11. Kirson ED, Gurvich Z, Schneiderman R, et al. Disruption of cancer cell replication by alternating electric fields. *Cancer Res*. 2004; 64(9):3288-3295.
12. Kirson ED, Schneiderman RS, Dbaly V, et al. Chemotherapeutic treatment efficacy and sensitivity are increased by adjuvant alternating electric fields (TTFIELDS). *BMC Med Phys*. 2009b; 9:1.
13. Mehta M, Wen P, Nishikawa R, et al. Critical review of the addition of tumor treating fields (TTFIELDS) to the existing standard of care for newly diagnosed glioblastoma patients. *Crit Rev Oncol Hematol*. 2017; 111:60-65.
14. Mrugala MM, Engelhard HH, Dinh Tran D, et al. Clinical practice experience with NovoTTF-100A™ System for glioblastoma: the patient registry dataset (PRiDe). *Semin Oncol*. 2014; 41 (Suppl 6):S4-S13.
15. Pless M, Droege C, von Moos R, et al. A phase I/II trial of Tumor Treating Fields (TTFIELDS) therapy in combination with pemetrexed for advanced non-small cell lung cancer. *Lung Cancer*. 2013; 81(3):445-450.
16. Regev O, Merkin V, Blumenthal DT, et al. Tumor-treating fields for the treatment of glioblastoma: A systematic review and meta-analysis. *Neurooncol Pract*. 2021; 8(4):426-440.
17. Rulseh AM, Keller J, Klener J, et al. Long-term survival of patients suffering from glioblastoma multiforme treated with tumor-treating fields. *World J Surg Oncol*. 2012; 10:220.
18. Salzberg M, Kirson E, Palti Y, Rochlitz C. A pilot study with very low-intensity, intermediate-frequency electric fields in patients with locally advanced and/or metastatic solid tumors. *Onkologie*. 2008; 31(7):362-365.
19. Sampson JH. Alternating electric fields for the treatment of glioblastoma. *JAMA*. 2015; 314(23):2511-2513.
20. Stupp R, Taillibert S, Kanner AA, et al. Maintenance therapy with tumor-treating fields plus Temozolomide vs Temozolomide alone for glioblastoma: a randomized clinical trial. *JAMA*. 2015; 314(23):2535-2543.
21. Stupp R, Taillibert S, Kanner A, et al. Effect of tumor-treating fields plus maintenance temozolomide vs maintenance temozolomide alone on survival in patients with glioblastoma: A randomized clinical trial. *JAMA*. 2017; 318(23):2306-2316.
22. Stupp R, Wong ET, Kanner AA, et al. NovoTTF-100A versus physician's choice chemotherapy in recurrent glioblastoma: a randomised phase III trial of a novel treatment modality. *Eur J Cancer*. 2012; 48(14):2192-2202.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

23. Toms SA, Kim CY, Nicholas G, Ram Z. Increased compliance with tumor treating fields therapy is prognostic for improved survival in the treatment of glioblastoma: A subgroup analysis of the EF-14 phase III trial. *J Neurooncol.* 2019; 141(2):467-473.
24. Trusheim J, Dunbar E, Battiste J, et al. A state-of-the-art review and guidelines for tumor treating fields treatment planning and patient follow-up in glioblastoma. *CNS Oncol.* 2017; 6(1):29-43.
25. Vymazal J, Wong ET. Response patterns of recurrent glioblastomas treated with tumor-treating fields. *Semin Oncol.* 2015; 42(3):e44-e55.
26. Wong ET, Lok E, Swanson KD, et al. Response assessment of NovoTTF-100A versus best physician's choice chemotherapy in recurrent glioblastoma. *Cancer Med.* 2014; 3(3):592-602.

Government Agency, Medical Society, and Other Authoritative Publications:

1. American Urological Association (AUA) / American Society for Radiation Oncology (ASTRO) / Society of Urologic Oncology (SUO). Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. 2017. Available at: https://www.astro.org/uploadedFiles/MAIN_SITE/Patient_Care/Clinical_Practice_Statements/Content_Pieces/ClinicallyLocalizedProstateCancer.pdf. Accessed on September 20, 2022.
2. Gilbert MR. Establishing the standard of care for patients with newly diagnosed and recurrent glioblastoma. *Am Soc Clin Oncol Educ Book.* 2012:112-117.
3. National Cancer Institute (NCI). Available at <http://www.cancer.gov/>. Accessed on September 20, 2022.
 - Adult Central Nervous System Tumors Treatment (PDQ®). Last modified January 18, 2022.
 - Malignant Mesothelioma Treatment (PDQ®). Last modified January 14, 2022.
 - Non-Small Cell Lung Cancer Treatment (PDQ®). Last modified March 17, 2022.
4. NCCN Clinical Practice Guidelines in Oncology®. © 2022 National Comprehensive Cancer Network, Inc. For additional information visit the NCCN website at: <http://www.nccn.org/index.asp>. Accessed on October 18, 2022.
 - Breast Cancer (V4.2022). June 21, 2022.
 - Central Nervous System Cancers (V2.2022). September 29, 2022.
 - Cutaneous Melanoma (V3.2022). April 11, 2022.
 - Malignant Pleural Mesothelioma (V1.2022). December 22, 2021.
 - Non-Small Cell Lung Cancer (V4.2022). September 2, 2022.
 - Pancreatic Adenocarcinoma (V1.2022). February 24, 2022.
5. Novocure.
 - Optune Instructions for Use. Issue Date: January 2019. Available at: https://www.optune.com/Content/pdfs/Optune_IFU_8.5x11.pdf. Accessed on September 20, 2022.
 - Patient Information and Operation Manual. Issue Date: January 2019. Available at: https://www.optune.com/content/pdfs/Optune_Piom_8.5x11.pdf. Accessed on September 20, 2022.
6. U.S. Food and Drug Administration (FDA). Humanitarian Device Exemption (HDE). NovoTTF™-100L System. Summary of Safety and Probable Benefit. H180002. May 23, 2019. Available at: https://www.accessdata.fda.gov/cdrh_docs/pdf18/H180002B.pdf. Accessed on September 20, 2022.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

7. U.S. Food and Drug Administration (FDA). Neurological Devices Panel. Available at: <https://www.fda.gov/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/NeurologicalDevicesPanel/>. Accessed on September 20, 2022.
8. U.S. Food and Drug Administration (FDA) Premarket Notification Database. Tumor Treatment Fields. NovoTTF-10A System. Summary of Safety and Effectiveness Data (SSED). No. P100034. Rockville, MD: FDA. April 8, 2011b. Available at: http://www.accessdata.fda.gov/cdrh_docs/pdf10/P100034b.pdf. Accessed on September 20, 2022.
9. U.S. Food and Drug Administration (FDA) Premarket Approval (PMA) Database. Premarket Notification Database. Tumor Treatment Fields. Optune™. Summary of Safety and Effectiveness Data (SSED). No. P100034/S013. Rockville, MD: FDA. October 5, 2015. Available at: http://www.accessdata.fda.gov/cdrh_docs/pdf10/p100034s013b.pdf. Accessed on September 20, 2022.

Websites for Additional Information

1. American Cancer Society (ACS).
 - What Are Adult Brain and Spinal Cord Tumors? Available at: <https://www.cancer.org/cancer/brain-spinal-cord-tumors-adults/about/what-are-brain-spinal-tumors.html>. Accessed September 20, 2022.
 - Malignant Mesothelioma. Available at <https://www.cancer.org/cancer/malignant-mesothelioma.html>. Accessed on September 20, 2022.
2. National Institutes of Health. Brain and Spinal Tumors Information Page. Available at: <https://www.ninds.nih.gov/Disorders/All-Disorders/Brain-and-Spinal-Tumors-Information-Page>. Accessed on September 20, 2022.

Index

NovoTTF-100A System
 NovoTTF-100L System
 Optune Lua
 Tumor Treatment Field (TTF)

The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.

History

Status	Date	Action
Revised	11/10/2022	Medical Policy & Technology Assessment Committee (MPTAC) review. Added medically necessary criteria for recurrent glioblastoma multiforme. Revised reference to see rationale for discussion about tumor progression criteria without change in intent. Updated Discussion and References sections.
Reviewed	08/11/2022	MPTAC review. Updated Discussion and References sections.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Electric Tumor Treatment Field (TTF)

Revised	08/12/2021	MPTAC review. Added medically necessary indications for continuation therapy. Updated Discussion and References sections.
Revised	08/13/2020	MPTAC review. Revised definition of tumor progression to refer reader to Discussion section. Updated Discussion and References sections. Reformatted Coding section.
Reviewed	08/22/2019	MPTAC review. Updated Discussion and References sections.
Revised	03/21/2019	MPTAC review.
Revised	03/20/2019	Hematology/Oncology Subcommittee review. Added a not medically necessary statement for treatment mapping and planning computer software. Updated Discussion and References sections.
New	05/03/2018	MPTAC review.
New	05/02/2018	Hematology/Oncology Subcommittee review. Initial document development. Moved content of DME.00035 Electric Tumor Treatment Field (TTF) to new clinical utilization management guideline document with the same title.

Federal and State law, as well as contract language including definitions and specific coverage provisions/exclusions, and Medical Policy take precedence over Clinical UM Guidelines and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Clinical UM Guidelines, which address medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Clinical UM Guidelines periodically. Clinical UM guidelines are used when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether or not to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the back of the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.